


Nidec

Power

Environmental Product Declaration



LSA 54.2

A **LEROY-SOMER**™ Product



**MADE IN EU
FOR EMEA**

Auditor's name: Abderezak Guiz

Product category rules based on "PCR-ed4"

Publication date: 07-2025

Validity period: 5 years

Independent verification of declaration and data in accordance with ISO 14025:2010

Internal External

Document complying with ISO 14025:2010 "Environmental labels and declarations. Type III environmental declarations"

NIDEC POWER'S ENVIRONMENTAL COMMITMENTS

Contributing to sustainable development

- Integrating environmental management into all of our activities through our ISO 14001 certification
- Decarbonizing our internal and external activities

Considering the environment in product design and providing information in compliance with ISO 14025


- Reducing the product environmental impact by integrating life cycle analysis from the design stage
- Working with our suppliers on the composition and origin of components

Offering our customers environmentally friendly solutions

- Developing a regionalization strategy to be closer to the user
- Providing efficient products that align with the context of electrification and grid intermittency

DESCRIPTION

Model	54.2
Reference	54.2 ZL18 – 3 812 kVA
Family	3 500 to 5 200 kVA - 50 Hz
Equipment	Low Voltage Alternator - 4 poles
Manufacturing	Czechia



VERIFICATION

Auditor's name: Abderezak Guiz	Product category rules based on "PCR-ed4"
Publication date: 07-2025	Validity period: 5 years
Independent verification of declaration and data in accordance with ISO 14025:2010	
Internal <input type="checkbox"/> External <input checked="" type="checkbox"/>	
Document complying with ISO 14025:2010 "Environmental labels and declarations. Type III environmental declarations"	

PRODUCT RANGE COVERED BY “EPD”

LSA 54.2 ZL18 is the heaviest reference product in the range. Extrapolation coefficients are available in the appendix for assessing the environmental impact of the other models in the range.

Total product weight: 10 780 kg (including packaging)

FUNCTIONAL UNIT AND REFERENCE FLOW


Ensure the production of 1 kVA using an auxiliary energy source 200 hours per year for a reference lifetime of 40 years.

Reference flows

Ratings 50Hz – 1 500 rpm		
	On the scale of the product over its reference lifetime (40 years)	At the level of the functional unit over its reference lifetime (40 years)
Equipment	1p 54.2 ZL18 or 10 700 kg	1/30 496 000 of 54.2 ZL18 or 3.50E-04 kg of product
Packaging	1 packaging or 80 kg	1/30 496 000 packaging or 2.26E-06 kg of packaging
Maintenance	2 bearings or 88 kg	1/30 496 000 bearings or 2.88E-06 kg of bearings


PRODUCT (WITHOUT PACKAGING) MASS BALANCE

METALS	Quantity	Others	Quantity
Magnetic steel	50-55%	Insulation	0-5%
Steel	30-35%	Others	0-5%
Copper	10-15%		
Grey casting	0-5%		
Power cable	0-5%		
Total	95-100%	Total	0-5%



LIFE CYCLE ANALYSIS

This statement is representative of the A1-A5+B2 breakdown (cradle to delivery + maintenance).



Purchased components mainly come from Europe and are assembled at the Olomouc site in Czechia.

Manufacturing processes were taken into account based on on-site measurements of the power consumption of the various production equipment.

Manufacturing processes external to Nidec Power are taken into account.

The energy model used is that of production in Czechia

- ✓ Electricity, low voltage {CZ} | market for | Cut-off
- ✓ Heat, central or small-scale, natural gas {RER} | market group for heat, central or small-scale, natural gas | Cut-off, U

Upstream transport of materials and components from suppliers' sites to the Olomouc plant has been taken into account.

Ferrous scrap generated during production is 80% recycled.

Aluminum scrap generated during production is 70% recycled.

This product travels 3 500 km by road to get from our manufacturing site to the installation site of our main customer.

This transport scenario is therefore representative of a product delivery in Europe.

- ✓ The transport model is: Transport, freight, lorry >32 metric ton, EURO6 {RER} | transport, freight, lorry >32 metric ton, EURO6 | Cut-off, U

This phase includes the disposal of the product packaging.

The environmental impact of the handling equipment used to install the product has not been taken into account (negligible impact).

Use scenario:

Energy consumption

Energy consumption for product use is not included in this declaration.

Maintenance

Alternator maintenance is covered by the scheme proposed by Nidec Power:

The bearings can be re-lubricated as standard	With regard to the type of grease, the lubricating intervals and the quantity to be supplied to each bearing: refer to nameplate directions	Minimum every 6 months
Standard grease	MOBIL POLYREX™ EM: NLGI grade 2	
Factory lubrication	MOBIL POLYREX™ EM: NLGI grade 2	
Special grease	Refer to nameplate directions	



Following the requirements of the nameplate is essential. Other priority information may be mentioned. Lubricating the alternator in operation and during the first commissioning is essential. Before using any other grease, check its compatibility with the original grease.

The nature and frequency of intervention over the typical life of the product are:

Maintenance equipment	Number of RLS interventions	Replacement frequency
Bearing	2	2 bearings, replaced once every 20 000 hours

By industry convention, the transport phase is taken into account on the assumption of a 100 km round trip by van.

The transport LCI is: Transport, passenger car, EURO5 {RoW} | transport, passenger car, EURO5 | Cut-off, U

The end-of-life of the alternator is not taken into account in this declaration.

ENVIRONMENTAL IMPACT ASSESSMENT

The Life Cycle Assessment on which this Environmental Product Declaration (EPD) is based complies with the criteria of PCR-ed4-FR-2021_09_14 of the PEP Ecopassport® Program.

The results were obtained using SimaPro 9.5 software and the ecoinvent 3.9 database. The characterisation method is EF3.1.

IMPACT INDICATORS PER kVA CORRESPONDING TO THE FUNCTIONAL UNIT


The “PEP” is based on the **production of 1 kVA per hour of operation**. The real impact of the life cycle stages of the product installed in a real situation must be calculated by the user of the “PEP” by multiplying the impact considered by the rated power according to the configuration chosen.

1500 rpm

Impact category	Unit	Total	A1	A2	A3	A4	A5	B2	Module D
PCR ed4 categories indicators									
Climate change	kg CO ₂ eq	1.83E-03	9.58E-04	5.32E-05	6.74E-04	1.24E-04	6.01E-07	2.03E-05	-1.49E-06
Climate change - Biogenic	kg CO ₂ eq	1.40E-05	2.62E-06	1.51E-08	1.03E-05	3.76E-08	2.06E-08	9.75E-07	-1.59E-09
Climate change - Fossil	kg CO ₂ eq	1.81E-03	9.55E-04	5.31E-05	6.63E-04	1.24E-04	5.80E-07	1.93E-05	-1.49E-06
Climate change - Land use and LU change	kg CO ₂ eq	1.75E-06	1.09E-06	2.83E-08	5.53E-07	6.04E-08	3.50E-11	1.96E-08	-1.15E-09
Ozone depletion	kg CFC-11 eq	3.43E-11	1.61E-11	1.14E-12	1.40E-11	2.81E-12	6.99E-15	2.62E-13	-1.28E-14
Acidification	mol H+ eq	3.27E-05	2.68E-05	3.62E-07	5.13E-06	3.07E-07	1.85E-09	1.41E-07	-1.03E-08
Eutrophication, freshwater	kg P eq	2.71E-06	2.18E-06	3.58E-09	5.06E-07	9.14E-09	7.85E-12	1.06E-08	-7.53E-10
Eutrophication, marine	kg N eq	2.72E-06	1.80E-06	9.29E-08	7.19E-07	8.36E-08	9.72E-10	2.29E-08	-2.58E-09
Eutrophication, terrestrial	mol N eq	3.23E-05	2.28E-05	1.00E-06	7.38E-06	8.58E-07	9.48E-09	2.14E-07	-2.92E-08
Photochemical ozone formation	kg NMVOC eq	1.15E-05	7.49E-06	3.68E-07	3.05E-06	5.01E-07	4.09E-09	7.33E-08	-1.04E-08
Resource use, minerals and metals	kg Sb eq	3.48E-07	3.10E-07	1.33E-10	3.61E-08	3.47E-10	2.25E-13	9.68E-10	-3.64E-12
Resource use, fossils	MJ	2.30E-02	1.15E-02	7.82E-04	8.63E-03	1.88E-03	4.59E-06	2.29E-04	-1.02E-05
Water use	m ³ depriv.	7.35E-04	5.58E-04	3.47E-06	1.60E-04	8.98E-06	5.71E-09	3.79E-06	-8.48E-07
Renewable primary energy w/t RM	MJ, net CV	2.96E-03	1.92E-03	1.07E-05	9.58E-04	2.75E-05	2.74E-08	4.90E-05	-1.26E-06
Renewable primary energy as RM	MJ, net CV	3.30E-06	0.00E+00	0.00E+00	3.30E-06	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Total renewable primary energy	MJ, net CV	2.97E-03	1.92E-03	1.07E-05	9.61E-04	2.75E-05	2.74E-08	4.90E-05	-1.26E-06
Non-renewable primary energy w/t RM	MJ, net CV	2.30E-02	1.15E-02	7.82E-04	8.63E-03	1.88E-03	4.59E-06	2.29E-04	-1.02E-05
Non-renewable primary energy as RM	MJ, net CV	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Total non-renewable primary energy	MJ, net CV	2.30E-02	1.15E-02	7.82E-04	8.62E-03	1.88E-03	4.59E-06	2.28E-04	-1.02E-05
Secondary materials	kg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Renewable secondary fuels	MJ, net CV	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Non-renewable secondary fuels	MJ, net CV	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Net use of freshwater*	m ³	-5.84E-05	-2.38E-05	-2.25E-07	-3.29E-05	-5.89E-07	-1.78E-10	-9.07E-07	2.00E-08
Hazardous wastes	kg	4.39E-04	2.95E-04	7.61E-07	1.26E-04	1.82E-06	6.17E-09	1.49E-05	-5.67E-07
Non-hazardous wastes	kg	1.00E-02	8.42E-03	6.54E-05	1.35E-03	1.79E-04	1.35E-07	3.35E-05	-4.99E-07
Radioactives wastes	kg	4.61E-08	2.36E-08	2.18E-10	2.12E-08	5.73E-10	5.00E-13	5.45E-10	-3.44E-11
Components for reuse	kg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Materials for recycling	kg	9.38E-05	0.00E+00	0.00E+00	9.38E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Materials for energy recovery	kg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Exported energy	MJ	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Total primary energy use	MJ, net CV	2.60E-02	1.34E-02	7.92E-04	9.58E-03	1.91E-03	4.61E-06	2.77E-04	-1.14E-05
PCR ed3 categories indicators									
PCR ed3-Global Warming	kg CO ₂ eq	1.79E-03	9.42E-04	5.24E-05	6.53E-04	1.22E-04	5.71E-07	1.91E-05	-1.48E-06
PCR ed3-Acidification soil & water	kg SO ₂ eq	2.83E-05	2.33E-05	2.88E-07	4.34E-06	2.44E-07	1.29E-09	1.19E-07	-8.12E-09
PCR ed3-Eutrophication	kg PO ₄ --- eq	9.49E-06	7.43E-06	4.46E-08	1.91E-06	6.14E-08	8.02E-10	4.69E-08	-3.27E-09
PCR ed3-Photochemical ozone creation	kg C ₂ H ₄ eq	2.69E-06	1.76E-06	6.27E-08	7.33E-07	1.27E-07	7.96E-10	1.42E-08	-2.28E-09
PCR ed3-Ozone Depletion	kg CFC-11 eq	3.74E-11	1.81E-11	9.35E-13	1.58E-11	2.30E-12	5.61E-15	2.59E-13	-1.02E-14
PCR ed3-Resource use, fossils	MJ	2.30E-02	1.15E-02	7.82E-04	8.63E-03	1.88E-03	4.59E-06	2.29E-04	-1.02E-05
PCR ed3-Resource use, min. and met.	kg Sb eq	3.48E-07	3.10E-07	1.33E-10	3.61E-08	3.47E-10	2.25E-13	9.68E-10	-3.64E-12
Optional categories indicators									
Optional-Particulate matter	disease inc.	1.63E-10	1.02E-10	4.67E-12	4.21E-11	1.23E-11	4.66E-14	1.40E-12	-1.95E-13
Optional-Ionising radiation	kBq U-235 eq	1.80E-04	9.17E-05	9.05E-07	8.28E-05	2.37E-06	2.25E-09	2.15E-06	-1.35E-07
Optional-Ecotoxicity, freshwater	CTUe	2.60E-02	1.92E-02	4.09E-04	5.28E-03	9.84E-04	2.18E-06	1.32E-04	-4.31E-06
Optional-Human toxicity, cancer	CTUh	1.42E-11	9.89E-12	2.35E-14	4.08E-12	5.51E-14	7.90E-17	1.11E-13	-8.02E-15
Optional-Human toxicity, non-cancer	CTUh	3.70E-10	3.26E-10	5.15E-13	4.19E-11	1.34E-12	1.61E-15	9.95E-13	-1.72E-14
Optional-Land use	Pt	1.69E-02	1.00E-02	6.96E-04	4.18E-03	1.91E-03	5.44E-07	1.19E-04	-4.40E-06
Product biogenic carbon content	0.00E+00	kg of C/UF							
Packaging biogenic carbon content	2.49E-06	kg of C/UF							

*Negative value linked to calculation methodology and data from database

DISTRIBUTION OF ENVIRONMENTAL IMPACTS



The raw materials extraction phase has the greatest impact on the indicators considered.

ENVIRONMENTAL IMPACT OF EQUIPMENT (additional information)

In the context of “LCA” at product level, the environmental impacts to be considered are those of the equipment expressed at product level over its reference lifetime.

Impact category	Unit	Total	A1	A2	A3	A4	A5	B2	Module D
PCR ed4 categories indicators									
Climate change	kg CO ₂ eq	5.58E+04	2.92E+04	1.62E+03	2.05E+04	3.78E+03	1.83E+01	6.19E+02	-4.55E+01
Climate change - Biogenic	kg CO ₂ eq	4.26E+02	8.00E+01	4.60E-01	3.14E+02	1.15E+00	6.27E-01	2.97E+01	-4.85E-02
Climate change - Fossil	kg CO ₂ eq	5.53E+04	2.91E+04	1.62E+03	2.02E+04	3.78E+03	1.77E+01	5.89E+02	-4.54E+01
Climate change - Land use and LU change	kg CO ₂ eq	5.34E+01	3.32E+01	8.63E-01	1.69E+01	1.84E+00	1.07E-03	5.98E-01	-3.52E-02
Ozone depletion	kg CFC-11 eq	1.04E-03	4.90E-04	3.48E-05	4.26E-04	8.58E-05	2.13E-07	7.99E-06	-3.90E-07
Acidification	mol H+ eq	9.97E+02	8.16E+02	1.10E+01	1.56E+02	9.35E+00	5.63E-02	4.30E+00	-3.14E-01
Eutrophication, freshwater	kg P eq	8.25E+01	6.64E+01	1.09E-01	1.54E+01	2.79E-01	2.39E-04	3.23E-01	-2.30E-02
Eutrophication, marine	kg N eq	8.29E+01	5.50E+01	2.83E+00	2.19E+01	2.55E+00	2.96E-02	6.99E-01	-7.86E-02
Eutrophication, terrestrial	mol N eq	9.84E+02	6.96E+02	3.06E+01	2.25E+02	2.62E+01	2.89E-01	6.52E+00	-8.92E-01
Photochemical ozone formation	kg NMVOC eq	3.50E+02	2.28E+02	1.12E+01	9.30E+01	1.53E+01	1.25E-01	2.24E+00	-3.16E-01
Resource use, minerals and metals	kg Sb eq	1.06E+01	9.46E+00	4.06E-03	1.10E+00	1.06E-02	6.86E-06	2.95E-02	-1.11E-04
Resource use, fossils	MJ	7.02E+05	3.51E+05	2.38E+04	2.63E+05	5.73E+04	1.40E+02	6.97E+03	-3.10E+02
Water use	m ³ depriv.	2.24E+04	1.70E+04	1.06E+02	4.89E+03	2.74E+02	1.74E-01	1.16E+02	-2.58E+01
Renewable primary energy w/t RM	MJ, net CV	9.03E+04	5.85E+04	3.25E+02	2.92E+04	8.39E+02	8.35E-01	1.50E+03	-3.86E+01
Renewable primary energy as RM	MJ, net CV	1.01E+02	0.00E+00	0.00E+00	1.01E+02	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Total renewable primary energy	MJ, net CV	9.04E+04	5.85E+04	3.25E+02	2.93E+04	8.39E+02	8.35E-01	1.50E+03	-3.86E+01
Non-renewable primary energy w/t RM	MJ, net CV	7.02E+05	3.51E+05	2.38E+04	2.63E+05	5.74E+04	1.40E+02	6.97E+03	-3.10E+02
Non-renewable primary energy as RM	MJ, net CV	0.00E+00							
Total non-renewable primary energy	MJ, net CV	7.01E+05	3.50E+05	2.38E+04	2.63E+05	5.73E+04	1.40E+02	6.97E+03	-3.10E+02
Secondary materials	kg	0.00E+00							
Renewable secondary fuels	MJ, net CV	0.00E+00							
Non-renewable secondary fuels	MJ, net CV	0.00E+00							
Net use of freshwater*	m ³	-1.78E+03	-7.25E+02	-6.85E+00	-1.00E+03	-1.79E+01	-5.44E-03	-2.77E+01	6.10E-01
Hazardous wastes	kg	1.34E+04	9.01E+03	2.32E+01	3.85E+03	5.54E+01	1.88E-01	4.55E+02	-1.73E+01
Non-hazardous wastes	kg	3.06E+05	2.57E+05	1.99E+03	4.12E+04	5.47E+03	4.12E+00	1.02E+03	-1.52E+01
Radioactives wastes	kg	1.40E+00	7.18E-01	6.65E-03	6.46E-01	1.75E-02	1.52E-05	1.66E-02	-1.05E-03
Components for reuse	kg	0.00E+00							
Materials for recycling	kg	2.86E+03	0.00E+00	0.00E+00	2.86E+03	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Materials for energy recovery	kg	0.00E+00							
Exported energy	MJ	0.00E+00							
Total primary energy use	MJ, net CV	7.91E+05	4.09E+05	2.42E+04	2.92E+05	5.82E+04	1.41E+02	8.46E+03	-3.48E+02
PCR ed3 categories indicators									
PCR ed3-Global Warming	kg CO ₂ eq	5.45E+04	2.87E+04	1.60E+03	1.99E+04	3.72E+03	1.74E+01	5.82E+02	-4.51E+01
PCR ed3-Acidification soil & water	kg SO ₂ eq	8.63E+02	7.11E+02	8.80E+00	1.32E+02	7.44E+00	3.94E-02	3.63E+00	-2.48E-01
PCR ed3-Eutrophication	kg PO ₄ --- eq	2.89E+02	2.27E+02	1.36E+00	5.82E+01	1.87E+00	2.44E-02	1.43E+00	-9.97E-02
PCR ed3-Photochemical ozone creation	kg C ₂ H ₄ eq	8.21E+01	5.35E+01	1.91E+00	2.23E+01	3.88E+00	2.43E-02	4.32E-01	-6.96E-02
PCR ed3-Ozone Depletion	kg CFC-11 eq	1.14E-03	5.52E-04	2.85E-05	4.82E-04	7.02E-05	1.71E-07	7.90E-06	-3.12E-07
PCR ed3-Resource use, fossils	MJ	7.02E+05	3.51E+05	2.38E+04	2.63E+05	5.73E+04	1.40E+02	6.97E+03	-3.10E+02
PCR ed3-Resource use, min. and met.	kg Sb eq	1.06E+01	9.46E+00	4.06E-03	1.10E+00	1.06E-02	6.86E-06	2.95E-02	-1.11E-04
Optional categories indicators									
Optional-Particulate matter	disease inc.	4.96E-03	3.12E-03	1.43E-04	1.28E-03	3.74E-04	1.42E-06	4.28E-05	-5.95E-06
Optional-Ionising radiation	kBq U-235 eq	5.48E+03	2.80E+03	2.76E+01	2.52E+03	7.23E+01	6.86E-02	6.55E+01	-4.13E+00
Optional-Ecotoxicity, freshwater	CTUe	7.93E+05	5.85E+05	1.25E+04	1.61E+05	3.00E+04	6.64E+01	4.02E+03	-1.32E+02
Optional-Human toxicity, cancer	CTUh	4.32E-04	3.02E-04	7.18E-07	1.24E-04	1.68E-06	2.41E-09	3.38E-06	-2.44E-07
Optional-Human toxicity, non-cancer	CTUh	1.13E-02	9.93E-03	1.57E-05	1.28E-03	4.10E-05	4.89E-08	3.04E-05	-5.25E-07
Optional-Land use	Pt	5.16E+05	3.06E+05	2.12E+04	1.27E+05	5.82E+04	1.66E+01	3.62E+03	-1.34E+02
Product biogenic carbon content	0.00E+00	kg of C							
Packaging biogenic carbon content	3.18E+01	kg of C							

*Negative value linked to calculation methodology and data from database

APPENDIX: extrapolation coefficients

The extrapolation coefficients can be applied to a homogeneous product family in accordance with the requirements of PCR-ed4-FR-2021_09_06.

The environmental impact of a product in the 54.2 range can be obtained using the following formula:

$$\text{Considered_product_impact} = \text{Reference_product_impact} \times \text{extrapolation_coefficient.}$$

For the manufacturing, distribution, installation and maintenance stages, the extrapolation coefficients are applied according to the following rule:

Coefficient on the scale of the declared product (additional information)	$\left(\frac{\text{total mass of the product considered with packaging (kg)}}{\text{total mass of the reference product of the range with packaging (kg)}} \right)$
--	--

3 300V			10 500V		
VL7	7 100	0.66	XL7	7 150	0.67
VL9	7 600	0.71	XL9	7 650	0.71
XL11	8 300	0.78	XL11	7 950	0.74
XL13	8 700	0.81	ZL12	8 450	0.79
ZL15	9 200	0.86	ZL14	8 900	0.83
4 160V			13 800V		
VL7	7 100	0.66	UL5	6 500	0.61
VL9	7 600	0.71	XL7	6 900	0.64
XL11	8 300	0.78	XL8	7 250	0.68
XL13	8 700	0.81	ZL10	7 800	0.73
ZL15	9 200	0.86	ZL12	8 000	0.75
6 300V			ZL14	8 400	0.79
VL7	7 050	0.66			
VL9	7 550	0.71			
XL11	8 300	0.78			
XL13	8 700	0.81			
ZL15	9 200	0.86			

GLOSSARY

Life cycle approach

Methodology for taking into account all the stages in the life of a product (manufacturing, installation, use and end of life) in order to determine the consequences for the environment.

Environmental aspect

Element of an organization's activities, products or services that may interact with the environment [ISO 14050].

Typical life cycle (TLC)

Theoretical life of the product retained for the functional unit.

Note: Typical life is a theoretical life used for calculation purposes. It can in no way be equated with the minimum, average or real life of products.

Homogeneous environmental family

Group of products corresponding to the same functional unit (identical function of use, product standard, identical technology (type of materials and manufacturing process) whose environmental impacts are identical to the reference product or can be extrapolated by applying a defined calculation rule.

Environmental impact

Any change to the environment, whether negative or beneficial, resulting in whole or in part from the activities, products or services of an organization [ISO 14050].

PCR (Product Category Rules)

A set of specific rules, requirements and guidelines for the development of Type III environmental declarations for one or more product categories [ISO 14025].

Reference product

Product or system of products modelled in the LCA and representative of a homogeneous environmental family.

Functional unit

Quantified performance of a product system intended to be used as a reference unit in a life cycle assessment [ISO 14040].



www.nidecpower.com

Connect with us at:



© 2025 Moteurs Leroy-Somer SAS. The information contained in this brochure is for guidance only and does not form part of any contract. The accuracy cannot be guaranteed as Moteurs Leroy-Somer SAS have an ongoing process of development and reserve the right to change the specification of their products without notice.

Moteurs Leroy-Somer SAS. Headquarters: Bd Marcellin Leroy, CS 10015, 16915 Angoulême Cedex 9, France. Share Capital: 32,239,235 €, RCS Angoulême 338 567 258.